

Diabetes in ABBEVILLE County, 2005



Produced by the South Carolina Department of Health and Environmental Control

Impact of Diabetes in the United States

■ Diabetes affects more than 23 million Americans and contributes to almost 234,000 deaths a year.


- Diabetes is becoming more common in the United States. From 1980 through 2007, the number of Americans with diabetes increased from 5.6 million to 17.9 million
- The direct and indirect costs of diabetes in the U.S. were nearly \$174 billion in 2007.
- 72,507 people died of diabetes in the U.S. in 2006.

Diabetes in South Carolina

- Approximately 332,000 adults in South Carolinians are diagnosed with diabetes. African Americans have twice the prevalence than Whites.
- The total direct costs of hospitalizations and emergency room visits were over \$8.5 billion in 2005.
- Diabetes is the seventh leading cause of death in South Carolina, claiming 1.183 lives in 2005.


Behavioral Risk Factors

■ Major behavioral risk factors, such as overweight, physical inactivity and unhealthy diet, are partially responsible for development of type 2 diabetes (Fig




Diabetes Prevalence

■ An estimated 1724 adults (8.7% of adults) in Abbeville County suffer annually from Diabetes (Fig 2).



Hospital and Emergency Room Discharges

- In 2005, there were 57 hospitalizations for Diabetes in Abbeville County for all ages.
- There were an additional 575 hospital discharges with diabetes-related conditions in Abbeville County.
- In 2005, in there were 49 emergency room (ER) visits for diabetes as the primary diagnosis, among which 36 (73%) were by African-American patients.
- There were 517 ER visits for diabetes as a related condition.




Medicare paid for 69% of the ER and Hospitalization charges in Abbeville County (Fig 3).

Complications

- Almost all patients (97%)hospitalized in SC for diabetes had complications in 2005; 36% of hospitalizations in Abbeville County (Table 1).
- In 2005, nearly three-quarters of people in SC with diabetes had hypertension, compared with only one-fourth of people without diabetes.

Table 1. Hospitalizations for Diabetes Complications Abbeville County in 2005.			
Diabetes Complication	Hospital Admissions	Percent of Diabetes Hospital Admissions	Total charges
Renal Failure	151	24%	\$3,616,400
Lower Extremity Amputation	7	1%	\$379,400
Ketoacidosis	23	4%	\$333,000
Retinopathy	48	8%	\$1,459,300

- In 2005, more than 60% of people with diabetes have high cholesterol.
- In 2005, the rates for lower extremity amputations among blacks in SC were four times the rates among whites.



For More Information about diabetes contact:

Division of Diabetes Prevention and Control: DHEC Bureau of Community Health and Chronic Disease Prevention

http://www.dhec.sc.gov/health/chcdp/diabetes/index.htm (803) 545-4471

Diabetes Initiative of South Carolina http://www.musc.edu/diabetes/


DIABETES IN A South Carolina

Generated by: Office of Chronic Disease Epidemiology, Bueau of Community Health and Chronic Disease Prevention


http://www.scdhec.gov/hs/epidata/state_reports.htm

Mortality

- Diabetes is the eighth leading cause of death in Abbeville County.
- A total of 5 people in Abbeville County died from diabetes in 2005.



- The Abbeville County mortality rate was 15.4/100,000 in 2005, below the state average.
- Abbeville County ranks #42 in diabetes mortality in South Carolina in 2003-2005.



Data Sources:

For more details, please view the Technical Document below www.scdhec.gov/health/epidata/docs/EpiTechNotes.pdf

Hospital Discharges and ER Visits, contact:

Office of Research and Statistics SC Budget and Control Board (803) 898-9958



